

Medical Necessity Guideline: Back Pain – Physical Therapy	Creation Date: 07/28/2014	Review Date: 05/24/2024	Effective Date: 06/11/2024
---	-------------------------------------	-----------------------------------	--------------------------------------

PURPOSE:

To define the conditions and requirements for therapy requests for acute and chronic “back pain” without associated trauma or neurological deficits.

LINE OF BUSINESS: STAR, STAR Kids, and CHIP

DEFINITIONS:

Back pain - a common disorder involving the muscles, nerves, and bones of the back. Pain can vary from dull constant ache to a sudden sharp feeling. Pain may be classified by duration as **Acute** (pain lasting less than 6 weeks), **Subchronic** (6-12 weeks), or **Chronic** (more than 12 weeks).

GUIDELINE:

Based on evidence-based studies and research, DHP considers appropriate physical therapy treatment of back pain, without associated trauma or neurological deficit, as short-term therapy tapering to low frequency with a transition to a robust home plan. ^(1, 2, 3)

Documentation Requirements: Therapy requests should include clinical notes with medical diagnoses, history of the pain to include a history of trauma or other etiology, chronicity, previous treatments or therapy, and a treatment plan that clearly justifies the proposed treatment frequency and intensity, and transition to a robust home program.

BACKGROUND:

Acute low back pain is defined as pain, increased muscle tone, and stiffness localized below the costal margin and above the inferior gluteal folds, sometimes accompanied by radiating pain, for up to 6 weeks: ⁽¹⁰⁾

- Pain that continues but does not exceed 12 weeks is defined as subacute
- Pain that continues past 12 weeks is defined as chronic
- The acute phase is further defined as early acute phase if it has been less than 2 weeks, and late acute phase if it is between 2 and 6 weeks
- If pain returns after complete recovery, it is considered acute recurrent

Acute back pain may result from various causes such as overexertion using the back muscles, poor posture including body mechanics and/or ergonomics, lumbar disc disease, facet joint

Confidential: For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

syndrome, sacroiliac joint dysfunction, and other pelvic girdle pain. However, a specific etiology is identified in only about 15% of cases.

Based on systematic reviews from Canada and Spain, good evidence supports physical therapy for pain management and patient education to stay active during the acute phase of low back pain (LBP).⁽⁴⁾ A current review of the literature demonstrates that there is insufficient evidence to support the use of supervised, individualized exercise therapy in acute LBP.^(3, 4, 5) However, investigators in the United States found that in patients with recent-onset LBP, early physical therapy resulted in statistically significant improvement in disability, but the improvement was modest and did not reach the minimum clinically significant difference compared with the usual care. The early physical therapy group received four physical therapy sessions over the course of 3 weeks, consisting of manipulation and exercise.

Authors of a 2018 systematic review conducted in China have concluded that patients with LBP who participate in an aquatic exercise program can have a significant reduction in pain, as well as improved physical functioning⁽⁶⁾

Maintenance or Prevention – Home Exercise and Instructions Plan - Continue exercises found effective and well-tolerated to maintain optimal physical functioning.^(5, 8) Avoidance of provocative activities, Patient education^(3, 7, 8, 9), and the use of shoe insoles and lumbar supports are not recommended for the prevention of LBP⁽⁵⁾

The prognosis in acute LBP is good, with recovery or improvement in 80 - 90% of cases after 6 - 8 weeks.⁽¹²⁾ However, in the first year following an acute episode, 60 - 86% of those affected experience a recurrence, and an estimated 2 - 3% develop chronic, disabling LBP.

PROVIDER CLAIMS CODES:

CPT				
97110	97112	97113	97116	97124
97140	97530	97535	97537	

ICD 10				
M54	M54.1	M54.3	M54.4	M54.5
M54.6	M54.8	M54.9	M48	M48.3
M48.4	M48.5	M48.8	M48.9	M53
M53.2	M53.3	M53.8	M53.9	

Confidential: For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

REFERENCES:

1. Texas Medicaid Provider Procedure Manual (Current Edition); Physical, Occupational, and Speech Therapy Services Handbook, 4.6 (Frequency and Duration Criteria for PT, OT, and ST services); May 2024.
2. Karlsson M, Bergenheim A, Larsson MEH, Nordeman L, van Tulder M, Bernhardsson S. Effects of exercise therapy in patients with acute low back pain: a systematic review of systematic reviews. *Syst Rev*. 2020 Aug 14;9(1):182. doi: 10.1186/s13643-020-01412-8. PMID: 32795336; PMCID: PMC7427286.; Accessed 05/18/2022.
3. Abuín-Porras V, Clemente-Suárez VJ, Jaén-Crespo G, Navarro-Flores E, Pareja-Galeano H, Romero-Morales C. Effect of Physiotherapy Treatment in the Autonomic Activation and Pain Perception in Male Patients with Non-Specific Subacute Low Back Pain. *J Clin Med*. 2021 Apr 20;10(8):1793. doi: 10.3390/jcm10081793. PMID: 33924113; PMCID: PMC8074333. Accessed 5/18/2022.
4. McIntosh G, Hall H. Low back pain (acute). *Clin Evid (Online)*. 2011; pii: 1102. ([SR](https://www.ncbi.nlm.nih.gov/pubmed/21549023)), <https://www.ncbi.nlm.nih.gov/pubmed/21549023>. Accessed 05/18/2022.
5. Malanga GA, Dunn KR. Low back pain management: approaches to treatment. *J Musculoskelet Med*. 2010;27(8):305-315. Retrieved from <https://go.gale.com/ps/anonymou?id=GALE%7CA263880279&sid=googleScholar&v=2.1&it=r&linkaccess=abs&issn=08992517&p=AONE&sw=w> . Accessed 05/18/2022.
6. Zhongju S, Hengxing Z, Lu BP, zhijian W, Xue Yao YK, Lu SF. Aquatic exercises in the treatment of low back pain: A systematic review of the literature and meta-analysis of eight studies. *Am J Phys Med Rehabil*. 2018;97(2):116-122. <https://www.ncbi.nlm.nih.gov/pubmed/28759476>. Accessed 05/18/2022.
7. Delitto A, George SZ, Van Dillen LR, et al. Low back pain. *J Orthop Sports Phys Ther*. 2012;42(4):A1. doi:10.2519/jospt.2012.0301. <https://www.ncbi.nlm.nih.gov/pubmed/22466247>. Accessed 05/18/2022.
8. National Guideline Clearinghouse Website (2022), Toward optimized practice. Guideline for the evidence-informed primary care management of low back pain. <https://www.guidelinecentral.com/summaries/guideline-for-the-evidence-informed-primary-care-management-of-low-back-pain/#section-date>. Accessed 05/18/2022.
9. Institute for Clinical Systems Improvement. Adult acute and subacute low back pain, Revised March 2018, Found at <https://www.icsi.org/guideline/low-back-pain/>. Accessed 05/18/2022.
10. Dressendorfer R, Back Pain: Low, Acute; CINAHL Rehabilitation Guide, EBSCO Publishing, (Ipswich, Massachusetts); 2019 Mar 15; Accessed 05/18/2022.
11. Atlas SJ, Deyo RA. Evaluating and managing acute low back pain in the primary care setting. *J Gen Intern Med*. 2001;16(2):120-131. doi:10.1111/j.1525-1497.2001.91141.x. Accessed 05/18/2022.
12. Lewis A, Morris ME, Walsh C. Are physiotherapy exercises effective in reducing chronic low back pain?. *Phys Ther Rev*. 2008;13(1):37-44. <https://www.tandfonline.com/doi/abs/10.1179/174328808X252000>. Accessed 05/18/2022.

Confidential: For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

DOCUMENT HISTORY:

DHP Committee that Approved	<i>Review Approval Date (last 5 years)</i>				
Medical Director	06/07/2022	05/23/2023	05/24/2024		
CMO	06/07/2022	06/06/2023	05/24/2024		
Medical Policy Workgroup	06/07/2022	06/06/2023	06/11/2024		
Utilization Management & Appeals	06/21/2022	06/20/2023	06/18/2024		
Provider Advisory Committee (PAC)	06/17/2022	06/09/2023	07/01/2024		
Clinical Management Committee	06/24/2022 & 08/23/2022	07/20/2023	07/24/2024		
Executive Quality Committee	06/28/2022	07/25/2023	07/30/2024		

<i>Document Owner</i>	<i>Organization</i>	<i>Department</i>
Dr. Fred McCurdy, Medical Director	Driscoll Health Plan	Utilization Management

<i>Review/Revision Date</i>	<i>Review/Revision Information, etc.</i>
08/07/2014	No change
09/01/2015	No change
11/01/2016	No change
03/30/2017	Recommendation by DHP Physical Therapists to address member with chronic LBP in a standard format.
11/01/2017	No change
11/19/2018	Added references

Confidential: For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

Driscoll Health Plan Medical Necessity Guideline



11/30/2019	Change in format, addition of descriptions, guidelines, and improved language. Significant updating of references and addition of additional references by Dr. Brendel.
05/07/2020	Update in format and language
06/04/2020	Format changes
06/16/2020	Final review
05/18/2021	Added 3 new references, validated other references and codes – Dr. Brendel
05/20/2022	Reviewed by Dr. Albert Gest
05/23/2022	Reviewed and edited by Dr. Fred McCurdy
05/23/2023	Reviewed by Drs Noorullah Akthar and Fred McCurdy
05/24/2024	Reviewed and Revised by Drs. Dan Doucet and Fred McCurdy

Confidential: For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan