

## Driscoll Health Plan Medical Necessity Guideline



<b>Medical Necessity Guideline:</b> Bell's Palsy-Physical Therapy	<b>Creation Date:</b> 09/05/2014	<b>Review Date:</b> 05/31/2023	<b>Effective Date:</b> 09/05/2014
--	-------------------------------------	-----------------------------------	--------------------------------------

### **PURPOSE:**

To define the requirements for therapy for the diagnosis of Bell's Palsy.

### **DEFINITIONS:**

**Bell's palsy** - an acute peripheral facial weakness of unknown cause, and the diagnosis can be established without difficulty in patients with unexplained unilateral isolated facial weakness. <sup>(2)</sup>

### **GUIDELINE:**

Corticosteroids, with or without antivirals, are the primary treatment of acute Bell's palsy. Evidence does not support the routine use of adjunctive modalities or therapies to hasten recovery or avoid progression to a chronic state. Members with Bell's palsy should receive education in the pathophysiology and natural course of the disease along with provider instruction in facial exercises and eye protection.

A physical therapy (PT), Occupational Therapy (OT), or Speech Therapy (ST) evaluation, along with limited visits, may be medically necessary for instruction on facial exercises and are primarily for the development and follow up of a home education program <sup>(1)</sup>

### **Documentation Requirements:**

Driscoll Health Plan (DHP) requires submission of:

- complete and detailed clinical notes describing the condition, whether acute or chronic
- all treatments provided, including corticosteroids and antivirals
- any referrals
- any current or previous POST therapy and modalities and response.

### **BACKGROUND:**

The incidence of Bell's palsy is 20-30 cases for 100,000 and accounts for 60-70% of all cases of unilateral peripheral facial palsy. Either sex is affected equally and may occur at any age, the median age is 40 years. The incidence is lowest under 10 years of age and highest in people over the age of 70. Left and right sides are affected equally <sup>(2)</sup> The terms "Bell's palsy" and "idiopathic facial paralysis" may no longer be considered synonymous, as herpes simplex virus activation is the likely cause of Bell's palsy in most cases. <sup>(3)</sup>

Medical Necessity Guideline: STAR, CHIP, STAR Kids

**Confidential:** For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

## Driscoll Health Plan Medical Necessity Guideline



Bell’s palsy is an acute disorder of the facial nerve, which produces full or partial loss of movement on one side of the face. The facial palsy gets completely better without treatment in most, but not all, people. <sup>(2)</sup> There is moderate to high confidence that corticosteroids are useful as a primary treatment for this condition. In addition, there is moderate evidence that the combination of corticosteroids and antivirals reduces the risk of long-term complications of Bell’s palsy.

There is insufficient or inconclusive evidence for the utility of biofeedback, laser, electrotherapy, massage, and thermotherapy in treating acute Bells’s Palsy. The mainstay in the treatment of acute Bell’s palsy remains corticosteroids. There is no clear benefit from antivirals alone as compared to placebo. However, the Cochrane Study did find moderate certainty that combination treatment with corticosteroids and antivirals reduced the number of people who experienced the long-term effects of Bell’s palsy. <sup>(3)</sup>

In Bell’s palsy various physical therapies, such as exercise, biofeedback, laser, electrotherapy, massage, and thermotherapy, are used to hasten recovery. However, the evidence for the efficacy of any of these therapies is lacking. A Cochrane systemic review of the effectiveness of physical therapies, electrostimulation, and exercises, on the outcome of Bell’s Palsy, concluded that there was no significant benefit or harm from any of these physical therapies for Bell’s Palsy. There was limited evidence that improvement began earlier in the exercise group. Another systematic review examined the effects of facial exercises associated either with mirror or electromyogram biofeedback with respect to complications of delayed recovery in Bell’s palsy and concluded that because of the small number of randomized controlled trials, it was not possible to analyze if the exercises, were effective. However, the possibility that facial exercise reduces the time to recover and sequelae needs to be confirmed with good-quality randomized controlled trials. <sup>(2, 3, 4)</sup>

**PROVIDER CLAIMS CODES:**

ICD 10	
G51.0	Bell’s Palsy

CPT			
97110	97112	97124	97140
97530	97535		

Medical Necessity Guideline: STAR, CHIP, STAR Kids

**Confidential:** For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

### **REFERENCES:**

1. Texas Medicaid Provider Procedure Manual (TMPPM) (May 2023); Physical, Occupational, and Speech Therapy Services Handbook, Section 4.5 (Frequency and Duration Criteria for PT, OT, and ST services).
2. Murthy, J. M., & Saxena, A. B. (2011). Bell's palsy: Treatment guidelines. *Annals of Indian Academy of Neurology*, 14(Suppl 1), S70–S72.  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3152161/>. Accessed 5/13/2022.
3. Bagyor I, Madhok VB, Daly F, Sullivan F. Antiviral treatment for Bell's palsy (idiopathic facial paralysis). *Cochrane Database of Systematic Reviews* 2019, Issue 9. Art. No.: CD001869. DOI: 10.1002/14651858.CD001869.pub9;  
[https://www.cochrane.org/CD001869/NEUROMUSC\\_antiviral-treatment-bells-palsy](https://www.cochrane.org/CD001869/NEUROMUSC_antiviral-treatment-bells-palsy);  
Accessed 05/13/2022.
4. Teixeira LJ, Valbuza JS, Prado GF. Physical therapy for Bell's palsy (idiopathic facial paralysis). *Cochrane Database of Systematic Reviews* 2011, Issue 12. Art. No.: CD006283. DOI: 10.1002/14651858.CD006283.pub3;  
[https://www.cochrane.org/CD006283/NEUROMUSC\\_physical-treatments-for-idiopathic-facial-paralysismt-bells-palsy](https://www.cochrane.org/CD006283/NEUROMUSC_physical-treatments-for-idiopathic-facial-paralysismt-bells-palsy); Accessed 05/13/2022.

Medical Necessity Guideline: STAR, CHIP, STAR Kids

**Confidential:** For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

## Driscoll Health Plan Medical Necessity Guideline



### DOCUMENT HISTORY:

DHP Committee that Approved	Review Approval Date (last 5 years)					
Medical Director	06/13/2019	06/22/2020	06/10/2021	05/24/2022	06/07/2022	05/31/2023
CMO	11/21/2018	6/13/2019	06/22/2020	06/10/2021	06/07/2022	06/01/2023
Medical Policy Workgroup Effective 2022					06/07/2022	06/06/2023
Medical Management Retired December 2020	11/21/2018	06/13/2019	06/22/2020			
Utilization Management & Appeals Effective January 2021				06/10/2021	06/21/2022	06/20/2023
Utilization Management Behavioral Health Retired December 2020	02/28/2019	08/22/2019	06/22/2020			
Provider Advisory Committee (PAC) Effective 2022					06/17/2022	06/09/2023
Clinical Management Committee				06/17/2021	06/24/2022 & 08/23/2022	07/20/2023

Medical Necessity Guideline: STAR, CHIP, STAR Kids

**Confidential:** For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

## Driscoll Health Plan Medical Necessity Guideline



<i>Effective March 2021</i>						
Quality Management <i>Retired 2020</i>	04/16/2019	10/22/2019	06/26/2020			
Executive Quality Committee <i>Effective 2021</i>				08/04/2021	06/28/2022	07/25/2023

<i>Document Owner</i>	<i>Organization</i>	<i>Department</i>
Dr. Fred McCurdy, Medical Director	Driscoll Health Plan	Utilization Management

<i>Review/Revision Date</i>	<i>Review/Revision Information, etc.</i>
03/04/2014	Original
09/05/2014	Modified
12/01/2015	No changes
10/06/2016	Modified-added APTA reference
12/01/2017	No changes
11/16/2018	No changes
11/30/2019	Modification of format, review of content and references
05/14/2020	Extensive rewording, updated references, codes added
06/16/2020	Dr. Serrao and Brendel edits
05/17/2021	Updated references, codes – Brendel – comments by Tamara Gonzales
05/13/2022	Reviewed and references updated by Dr. D. Doucet
05/24/2022	Final review and final edits by Dr. Fred McCurdy
05/31/2023	Reviewed by Drs. Dan Doucet and Fred McCurdy; no changes

Medical Necessity Guideline: STAR, CHIP, STAR Kids

**Confidential:** For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan