



Medical Necessity Guideline: Pes Planus (Flat	Creation	Review	Effective
Feet/Valgus Deformity) and Orthotics	Date:	Date:	Date:
	3/14/2014	05/24/2024	06/11/2024

PURPOSE:

To detail the authorization requirements for Orthotics Pes Planus (Flat Feet/Valgus deformity) and related conditions

DEFINITIONS:

Pes Planus (Flat foot) - The clinical diagnosis of flat foot is based on a valgus position of the heel and poor formation of the arch.

GUIDELINE:

Orthotic

A University of California at Berkeley Laboratory (UCBL) removable foot insert will be considered for prior authorization with documentation that the device is required to correct or treat at least one of the following conditions: ²

- A valgus deformity and significant congenital pes planus with pain.
- A structural problem that results in significant pes planus, such as Down syndrome.
- Acute plantar fasciitis.

Non-covered Orthotic Services:¹

- Tired feet
- Fatigued feet
- Non-severe bow legs
- Valgus deformity of the foot (except as outlined in the orthotic section see above)
- Pes planus (flat feet) (except when there is a coexisting medical condition outlined in the Texas Medicaid Provider Procedures Manual ²)

Required Documentation: ⁽⁵⁾

- Clear description of the term "flat feet" including age and whether physiologic/developmental as in early childhood, flexible-flat foot, or true pes planus.
- The use of the term "acquired pes planus" or other "acquired" condition must be explained in detail including the causative mechanism or condition.
- Documentation of the description and the qualification of pain associated with the diagnosis. The use of "painful feet" alone is considered inadequate to meet this standard.

Medical Necessity Guideline: STAR, CHIP, STAR Kids

Confidential: For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

4525 Ayers St. Corpus Christi, Texas 78415 Telephone (877) 455-1053 Fax (866) 741-5650





• Attestation and description of previous failed education and 6-month trials of over-thecounter Orthotics and compliance with purchase and use of appropriate non-prescription footwear.

BACKGROUND:

Natural history of Pes Planus (Flat Foot)

Pes planus is common in infants and children. It usually resolves by the time a person is an adolescent. Therefore, pes planus is typically described as physiologic because it is common and usually doesn't cause symptoms (i.e., the foot is typically flexible and without pain). In addition, there are no functional consequences to the foot being "flat".

Infants are usually born with flexible, flat feet and a very large fat pad on the bottom of the foot that obscures the medial plantar arch. During the first decade of life, this medial longitudinal arch develops along with the foot's bones, muscles, and ligaments, and by age two (2), a medial arch will be visible when the child is sitting. When the child stands (bears weight), this medical arch will frequently disappear, giving the appearance of flat feet. This usually resolves by the time the child is ten (10), but flat feet may persist into adolescence and adulthood. There is debate in the medical community as to whether this should be considered a normal variant or a deformity that leads to deformity requiring intervention. If there are no symptoms, most physicians would agree that this is a normal variant of foot shape that will persist throughout the person's lifetime. In rare instances, however, pes planus can become painful, and the foot may lose flexibility. This is frequently the first sign of some underlying foot pathology (e.g., arthritis, neuromuscular disease, or tarsal coalition)³.

Definition of Pes Planus

Despite being common, there is no standard definition for pes planus. Furthermore, no extensive, prospective studies compare the natural history of pes planus throughout the growth and development, comparing the response to various treatments. The available literature does not elucidate which patients are at risk for developing pain and disability as young adults. Current evidence suggests that observing an asymptomatic child with flat feet is safe and appropriate. Orthotics, although generally unproven to alter the course of flexible flatfoot, may provide relief of pain when pain is present. A general pediatrician needs to know when a referral to an orthopedic specialist is indicated, and which treatments may be offered to the patient. ⁴

Epidemiology of Pes Planus

Numerous cross-sectional epidemiologic studies have shown that pes planus is the normal shape of the foot in the first few years of life. In children two (2) years or younger, 97% have pes planus when the heel-to-arch width ratio defines it. This prevalence declines dramatically with

Medical Necessity Guideline: STAR, CHIP, STAR Kids





age, such that only 4% of children have pes planus by age ten (10). This supports the belief that most pediatric pes planus resolves spontaneously throughout the first decade of life. A study analyzing footprints in >800 patients found a similar trend, with 54% of 3-year-old children having flat feet. The prevalence then decreased to 26% when the children reached age six (6). This data suggests that the time between ages three to six (3 to 6) years may be a critical period for the development of the medial longitudinal arch. Of note, this same study looked at footprints in people eighty (80) years of age and discovered that pes planus is a normal adult finding. ⁵

Intervention in Pes Planus

Arch supports have long been advocated with very little evidence of benefit. Short-term positive results in the reduction of symptoms have been variably shown. However, no long-term follow-up studies have led many authors to conclude that treating children with physiologic pes planus is ineffective. It has also been posited that these treatments are uncomfortable and embarrassing for the child. One prospective study was conducted to determine if flexible pes planus could be influenced by corrective shoes and inserts and concluded that wearing corrective shoes or inserts for three (3) years did not affect the course of flexible flatfoot in children. Many other studies have discovered similar inability of custom foot orthotics or specialized shoes to be ineffective in treating pediatric flatfoot. Furthermore, other studies have shown no radiologic improvement with the use of arch supports/custom foot orthotics. ⁶

Current Approach to Managing Pes Planus in Children

Although the natural history of the flexible flatfoot remains unknown, there is little evidence to suggest that this condition results in long-term problems or disability. Treatment is reserved for a small subset of patients who develop symptoms. Patients may complain of hindfoot pain, abnormal shoe wear, or fatigue after long walking. These patients may benefit from a nonprescription orthosis, such as a medial arch support. Severe cases, often associated with an underlying connective tissue disorder such as Ehlers-Danlos syndrome or Down syndrome, may benefit from a custom orthosis such as the UCBL (University of California Biomechanics Laboratory) orthosis to better control the hindfoot and prevent collapse of the arch. Although an orthosis may relieve symptoms, there is no evidence to suggest any permanent change in the foot's shape or alignment of the tarsal bones.^{7,8}

ICD 10	
M21.4	Flatfoot (acquired)
M21.40	Flatfoot (acquired) unspecified foot
M21.41	Flatfoot (acquired) right foot
M21.42	Flatfoot (acquired) left foot
M21.6	Acquired deformities of foot

PROVIDER CLAIMS CODES:

Medical Necessity Guideline: STAR, CHIP, STAR Kids





Q66.5	Congenital pes planus
Q66.50	Congenital pes planus unspecified foot
Q66.51	Congenital pes planus right foot
Q66.52	Congenital pes planus left foot
Q66.6	Valgus deformities of feet

	СРТ
L3000	Foot, insert, removable, molded to patient model, 'ucb' type, berkeley shell, each
L3010	Foot, insert, removable, molded to patient model, longitudinal arch support,
	each
L3020	Foot, insert, removable, molded to patient model, longitudinal/metatarsal
	support, each
L1907	Ankle orthosis, supramalleolar with straps, with or without interface/pads,
	custom fabricated

Medical Necessity Guideline: STAR, CHIP, STAR Kids

Confidential: For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

4525 Ayers St. Corpus Christi, Texas 78415 Telephone (877) 455-1053 Fax (866) 741-5650





REFERENCES:

- 1. Texas Medicaid Provider Procedures Manual (Current Edition); Durable Medical Equipment, Medical Supplies, and Nutritional Substances Handbook; Section 2.2.19 (Orthotic Services [CCP]), 2.2.19.1 (Non-covered Orthotic Services), May 2024.
- Texas Medicaid Provider Procedures Manual (Current Edition); Durable Medical Equipment, Medical Supplies, and Nutritional Substances Handbook; Section 2.2.19.2.3 (Foot Orthoses), May 2024.
- Babu D, Bordoni B. Anatomy, Bony Pelvis and Lower Limb: Medial Longitudinal Arch of the Foot. [Updated 2022 Nov 11]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <u>https://www.ncbi.nlm.nih.gov/books/NBK562289/</u>
- 4. James B. Carr, Scott Yang, Leigh Ann Lather; Pediatric Pes Planus: A State-of-the-Art Review. Pediatrics March 2016; 137 (3): e20151230. 10.1542/peds.2015-1230
- 5. James B. Carr, Scott Yang, Leigh Ann Lather; Pediatric Pes Planus: A State-of-the-Art Review. Pediatrics March 2016; 137 (3): e20151230. 10.1542/peds.2015-1230
- Bresnahan PJ, Juanto MA. Pediatric Flatfeet-A Disease Entity That Demands Greater Attention and Treatment [published correction appears in Front Pediatr. 2021 Sep 01;9:735481]. Front Pediatr. 2020;8:19. Published 2020 Feb 11. doi:10.3389/fped.2020.00019
- Winell, Jennifer J, Davidson, Richard S., The Foot and Toes, <u>Nelson Textbook of Pediatrics</u>, 21st ed. Philadelphia, Pa: Saunders Elsevier; 2008:. Chapter 694, 3597-3606.e2. https://www.clinicalkey.com/#!/content/book/3-s2.0-B9780323529501006945?scrollTo=%23top. Retrieved September 18, 2023.
- Ueki Y, Sakuma E, Wada I. Pathology and management of flexible flat foot in children. *J Orthop Sci.* 2019 Jan;24(1):9-13. doi: 10.1016/j.jos.2018.09.018. Epub 2018 Oct 23. PMID: 30366675.; Accessed 05/17/2022.

Medical Necessity Guideline: STAR, CHIP, STAR Kids





DOCUMENT HISTORY:

DHP Committee that Approved		Review	[,] Approval Date	e (last 5 years)	
Medical Director	05/23/2023	10/10/2023	05/24/2024		
СМО	06/06/2023	12/13/2023	06/11/2024		
Medical Policy Workgroup	06/06/2023	12/13/2023	06/11/2024		
Utilization Management & Appeals	06/20/2023	12/13/2023	06/18/2024		
Provider Advisory Committee (PAC)	06/09/2023	12/13/2023	07/01/2024		
Clinical Management Committee	07/20/2023	01/24/2024	07/24/2024		
Executive Quality Committee	07/25/2023	01/30/2024	07/30/2024		

Document Owner	Organization	Department
Dr. Fred McCurdy, Medical Director	Driscoll Health Plan	Utilization Management

Review/Revision Date	Review/Revision Information, etc.		
6/1/2015	No change		
6/1/2016	No change		
12/1/2017	No change		
11/16/2018	No change		
11/15/2019	Update references and conversion to new format by Tom Morris, MD; addition of process. No substantive change to benefit guideline content		

Medical Necessity Guideline: STAR, CHIP, STAR Kids





05/12/2020	Added/updated new codes and new references
06/16/2020	Dr. Serrao's style recommendations added and modified, Add new source
05/21/2021	Updated, 2 new references, codes verified by Dr. Brendel
05/20/2022	Reviewed and updated, Dr. Albert Gest
05/23/2022	Reviewed and edited, Dr. Fred McCurdy
05/23/2023	Reviewed by Drs. Noorullah Akthar and Fred McCurdy; no changes
10/10/2023	Clarification of Non-Covered Orthotics services and Background
	revisions.
05/24/2024	Reviewed and revised by Michelle Hays, PT, and Dr. Fred McCurdy

Medical Necessity Guideline: STAR, CHIP, STAR Kids

Confidential: For use only by employees and authorized agents of Driscoll Health Plan. This document contains confidential and proprietary information NOT to be reproduced or distributed to others without the prior written consent of Driscoll Health Plan

4525 Ayers St. Corpus Christi, Texas 78415 Telephone (877) 455-1053 Fax (866) 741-5650